

On the Four Dimensional Conformal Anomaly, Fractal Spacetime and the Fine Structure Constant

Carlos Castro

Center for Theoretical Studies of Physical Systems
Clark Atlanta University, Atlanta, Georgia, 30314

October 2000

Abstract

Antoniadis, Mazur and Mottola (AMM) two years ago computed the intrinsic Hausdorff dimension of spacetime at the infrared fixed point of the quantum conformal factor in $4D$ Gravity. The fractal dimension was determined by the coefficient of the Gauss-Bonnet topological term associated with the conformal gravitational anomaly and was found to be greater than 4. It is explicitly shown how one can relate the value of the Hausdorff dimension computed by AMM to the universal dimensional fluctuation of spacetime ϵ given by $\phi^3/2$, where ϕ is the Golden Mean 0.618... Based on the infrared scaling limit of the theory and using recent Renormalization Group arguments by El Naschie, we conjecture that the unknown coefficient Q^2 , associated with the four dimensional gravitational conformal anomaly, could be precisely equal to the inverse fine structure constant values ranging between 137.036 and 137.081. Our results generate decimal digits up to any *arbitrary* number.

The Conformal Anomaly and Fractal Spacetime at Large Scales

Antoniadis, Mazur and Mottola [1] more than two years ago computed the intrinsic Hausdorff dimension of spacetime at the infrared fixed point of the quantum conformal factor in $4D$ Gravity. The fractal dimension was determined by the coefficient of the Gauss-Bonnet *topological* term associated with the conformal anomaly : *trace anomaly* and was found to be greater than 4. They also discussed a plausible physical mechanism for the *screening* of the cosmological constant at very large distances in full agreement with Nottale's work [7] and other results obtained by the present author and collaborators [6].

The Hausdorff dimension for spacetime is related to the geodesic distance $l(x, x')$ between points x, x' and the volume V_l enclosed by the spherical surface radius equal to l . The scaling relation between the two is $V_l \sim l^{d_H}$. For large l , this scaling relation defines the intrinsic dimension d_H of the space.

In $2D$ quantum gravity [2,3] the most appropriate way to calculate the d_H is by the heat kernel methods associated with the Laplacian operator $D_\mu D^\mu$ and the proper time s :

$$K_2(x, x'; s, g) = \langle x | e^{-s D_\mu D^\mu} | x' \rangle . \quad (1)$$

The heat kernel K_2 has a short distance expansion whose anomalous scaling behaviour can be calculated based on the standard techniques pioneered by B. de Witt in the sixties. The average geodesic length-squared that a scalar particle is able to diffuse after a proper time s is given by :

$$l_s^2 \equiv \frac{1}{V} < \int d^2 x \sqrt{g} \int d^2 x' \sqrt{g'} l^2 K_2(x, x'; s, g) >_V . \quad (2)$$

where the average is taken with respect to the fixed volume Liouville field theory partition function. By expanding the heat Kernel $K_2(x, x'; s; g)$ in a power series s one can see that

$$l_s^2 \sim s \quad \text{as} \quad s \rightarrow 0. \quad (3a)$$

which is the standard result undergoing Brownian motion.

The relevant scaling behaviour (with area/volume) is the expectation value of the following quantity appearing in the heat kernel expansion, under the conformal scalings of the metric $g_{ab} = \exp(2\sigma)\bar{g}_{ab}$:

$$s < \int d^2 x \sqrt{g} D_\mu D^\mu f_\epsilon(x, x_o) >_V = s < \int d^2 x \sqrt{\bar{g}} \bar{D}_\mu \bar{D}^\mu \bar{f}_\epsilon(x, x_o) >_V \sim s V^{\frac{\alpha-1}{\alpha_1}} \quad (3b)$$

where $f_\epsilon(x, x')$ is any smooth function with support only for distances of $l(x, x') \sim |x - x'| < \epsilon$. As ϵ goes to zero it approaches a delta function : $(1/\sqrt{g})\delta^2(x - x')$

The finite area/volume scaling behaviour of the last proportionality factor follows by a constant shift in the Liouville field $\sigma \rightarrow \sigma + \sigma_o$. The α_n are the anomalous scaling dimensions associated with the fields of the Liouville field theory and are given in terms of the weights n by the formulae :

$$\alpha_n = n + \frac{\alpha_n^2}{Q^2} = \frac{2n}{1 + \sqrt{1 - \frac{4n}{Q^2}}}. \quad (4a)$$

where the charge Q^2 is determined in terms of the matter central charge (anomaly coefficient) c_m by :

$$Q^2 = \frac{25 - c_m}{6}. \quad D = 2. \quad (4b)$$

We refer to the references [1,2,3] for further details. The main result is that the scaling behaviour of s under a global area/volume scaling

$$s \rightarrow \lambda^{-\frac{\alpha-1}{\alpha_1}} s. \quad (5)$$

will determine the Hausdorff Dimension d_H from the relation :

$$l_s^2 \sim s \sim V_l^{-\frac{\alpha-1}{\alpha_1}} \sim l^{-d_H \frac{\alpha-1}{\alpha_1}}. \quad (6)$$

giving finally :

$$d_H = -2 \frac{\alpha_1}{\alpha-1} = 2 \frac{\sqrt{25 - c_m} + \sqrt{49 - c_m}}{\sqrt{25 - c_m} + \sqrt{1 - c_m}} \geq 2. \quad D = 2 \quad (7)$$

Notice that the classical limit is obtained in eqs-(4a,4b,7) by setting $c_m = -\infty$ which implies that $Q^2 = \infty$ so that in eq-(7) the Hausdorff $d_H = 2$ as it ought to. The authors [1] repeated this analysis associated with the conformal anomaly in $D = 4$ where the charge Q^2 is now the coefficient of the Gauss-Bonnet curvature squared terms present in the four dimensional conformal anomaly :

$$Q^2 = \frac{1}{180}(N_S + \frac{11}{2}N_{WF} + 62N_V - 28) + Q_{grav}^2. \quad (8)$$

the *unknown* charge Q^2 is given in terms of the number of free scalars N_S , Weyl fermions N_{WF} and vector fields N_V . While -28 and the *unknown* value of the gravitational charge Q_{grav}^2 are the contributions of the spin-0 conformal factor and spin-2 graviton fields of the metric itself.

The scaling behaviour of the proper time under a global scaling of the volume $V \rightarrow \lambda V$ in $D = 4$ is :

$$s \rightarrow \lambda^{-\frac{\beta_8}{\beta_o}} s. \quad (9)$$

where β_o, β_8 are the conformal scaling exponents corresponding to the volume operator and the higher quartic derivative operator (square of the Laplacian plus curvature and other derivative terms). The Hausdorff dimension is given, in the long scale limit, by the relation :

$$l_s^4 \sim s \sim V_l^{-\frac{\beta_8}{\beta_o}} \sim l_s^{-d_H \frac{\beta_8}{\beta_o}} \quad (10)$$

then the Hausdorff dimension of fractal spacetime is explicitly given by :

$$d_H = -4 \frac{\beta_o}{\beta_8} = 4 \frac{1 + \sqrt{1 + \frac{8}{Q^2}}}{1 + \sqrt{1 - \frac{8}{Q^2}}} \geq 4. \quad (11)$$

This final expression for the spacetime fractal dimension is **all** we need to show that the *unknown* charge Q^2 in eq-(8) can be equated with the inverse of the fine structure constant 137.036.... (given by the Particle data booklet).

The authors [1] emphasized that the value of Q^2 was uncertain , principally because of the unknown infrared contributions of gravitons to the value of Q_{grav}^2 which appears in the r.h.s of eq-(8).

To calculate what the value of Q^2 may be , which in turn will yield the value of Q_{grav}^2 present in the r.h.s of (8), we conjecture that this value can be related to the inverse of the fine structure constant 137.036.. based on the recent papers by the author, Granik and El Naschie [4,6] . The main results of [4] and [6] is that there is a *universal* dimensional fluctuation in Nature given in terms of the Golden Mean by :

$$\Delta D_{fluctuation} = \epsilon = \frac{\phi^3}{2} = \phi - \frac{1}{2} = (0.618 - \frac{1}{2}) = 0.118..... \text{ where } \phi + 1 = \frac{1}{\phi} \Rightarrow \phi = \frac{\sqrt{5}-1}{2} = 0.618.... \quad (12)$$

and that the inverse of the fine structure constant [4] , ranging between 137.036... and El Naschie's results 137.081..., can be thought of as an *internal* dimension or a *central* charge, using the language of Irrational Conformal field Theory, as Eddington envision long ago. This line of reasoning is nothing but following the path chartered by Einstein himself on the geometrization of *all* physics, with the new ingredient that we believe that Nature **is** fractal at its core.

A simple numerical calculation shows that by simply setting $Q^2 = 137.036....$ inside the basic equation (11) yields automatically nothing more, nothing else but $4+\epsilon$ for the Hausdorff dimension of fractal spacetime , where ϵ is the universal dimensional fluctuation given by $\phi^3/2$:

$$4 \frac{1 + \sqrt{1 + \frac{8}{137.036}}}{1 + \sqrt{1 - \frac{8}{137.036}}} = 4.11856 \sim 4 + \frac{\phi^3}{2} = 4.118..... \quad !!!!!. \quad (13)$$

Is this a **numerical concidence** or **design** ???? . Conversely, we can propose an **exact** value for the inverse of the fine structure constant simply by setting the right hand side of (13) to be precisely

$$d_H = 4 + \epsilon = 4 + \phi^3/2 = 4 + \frac{\sqrt{5}-2}{2} \equiv 4 \frac{1 + \sqrt{1 + \frac{8}{Q^2}}}{1 + \sqrt{1 - \frac{8}{Q^2}}}. \quad (14)$$

Eq-(14) will determine the *precise* value of $Q^2 = 137.....$ up to an *arbitrary* number of digits. This is very encouraging because the only thing we have to do is to compare the value we get for the inverse fine structure constant, Q^2 , and compare with the results of **QED** !!! This could very well be the **sought-after numerical** proof of the correctness of the theory. Evenfurther, notice that in this infrared limit, we are working entirely with ordinary $4D$ gravity (long distance limit of string theory) and there is no need to do any compactifications from higher to lower dimensions.

The graph of $d_H = d_H(Q^2)$ is a decreasing function of Q^2 (the internal dimension in Eddington's language) . When $Q^2 \rightarrow \infty$ the $d_H = 4$ and one recovers the classical four dimensional limit. When $Q^2 < 8$ the dimension becomes *complex* which is not so a farfetched thing to imagine. One can envision an analytical continuation of the dimension consistent with the complexification of $\mathcal{E}^{(\infty)}$ spacetime and the celebrated Banach-Tarski theorem. One can imagine a sort of Quantum/Classical *duality* as follows :

Earlier in eqs-(4a,4b,7) we discussed that the classical limit is recovered when the matter central charge (or the embedding target spacetime dimension using the language of string theory) is $c_m = d = -\infty$; $Q^2 = \infty$ so that the Hausdorff dimension of the *worldsheet* is now $d_H = 2$. This point $d = -\infty$ has for *antipodal point* the one of $d = \infty$ which correspond, respectively, to the topological dimensions associated with the totally *void* set $\mathcal{E}^{(-\infty)}$ and the total space $\mathcal{E}^{(\infty)}$. The former has for Hausdorff dimension the value of 0 and the latter has for Hausdorff dimension ∞ . It is clear now why the void space $\mathcal{E}^{(-\infty)}$ is the dimension-dual space to $\mathcal{E}^{(\infty)}$ since $\infty = 1/0$. In this sense, we can argue that there a sort of Quantum/Classical duality principle operating.

The Banach-Tarski theorem, which roughly states that you can get two objects out of one , all of them identical, by abandoning the Axiom of Choice in Set Theory, fits very well with the *complexification* of $\mathcal{E}^{(\infty)}$ space by having another *dual* space : $\mathcal{E}_{dual}^{(\infty)}$ or *mirror* image. One can then perform the analytical continuation (or duality quantum/classical transformation) from $d = \infty$ to $d = -\infty$, by moving along a

large circle (infinite radius) in the complex-dimensions plane , and reach $d = -\infty$ from $d = \infty$. And then start over again and again moving in infinite cycles . The quantum *vacuum* , at $d = \infty$, is in this picture both a *source* and a *sink*. The creation of our Universe out of “ nothing “ could have occurred in this fashion. This resembles the picture of the Snake (coined *ouraborus*) eating its tail, then re-emerging again out of the vacuum and eating its tail later on and beginning another cycle, again and again, ad infinitum.

After this slight detour on the nature of *complex* dimensions, we conclude by saying that El Naschie [4] has presented very convincing arguments regarding the unification of gravity with the electroweak and strong forces, based on Renormalization Group Arguments, supporting that there is a very deep and explicit connection between Nottale’s Scale Relativity [7] , Irrational Conformal Field theory, El Naschie’s Cantorian-Fractal spacetime and the New Extended Scale Relativity formulated by the author [5], and that the inverse fine structure constant (an internal dimension or charge) plays a fundamental role in determining the scaling regimes of the electroweak and strong interaction as well as the hierarchy of the 16,6 internal dimensions, present in the Heterotic string theory and its compactifications , from $26 \rightarrow 10 \rightarrow 4$. Argyris et al [8] have recently shown how a fractalization of spacetime may be an intrinsic property of all processes in Nature, from the microworld to cosmos, having well defined signatures in Cosmic Strings and in the phenomenon of Spontaneous Symmetry breaking .

Acknowledgements

The author thanks M. S. El Naschie for his hospitality and for the series of discussions which led to this work. Also we wish to thank S. Ansoldi for his assistance and to E. Spallucci, E. Gozzi, M. Pavsic, A. Granik, T. Smith, C. Handy, A. Schoeller, A. Boedo, J. Mahecha, J. Giraldo, L. Baquero for their support.

References

1. Antoniadis, P. Mazur and E. Mottola : “ Fractal Geometry of Quantum Spacetime at Large Scales “ hep-th/9808070 .
- Antoniadis, P. Mazur and E. Mottola : Nuc. Phys. **B 388** (1992) 627
- 2- V. Knizhnik, A. Polyakov and A. Zamolodchikov : Mod. Phys. Lett **A 3** (1988) 819.
- F. David, Mod. Phys. Lett **A 3** (1988) 1651.
- F. David : Nuc. Phys. **B 257** (1985) 45
- 3- J. Distler, H. Kawai : Nuc. Phys. **B 257** (1985) 509
- V. Kazakov, Phys. Lett **B 150** (1985) 282
- L. Ambjorn, B. Durhuus, J. Frohlich and P. Orland : Nuc. Phys. **B 270** (1986) 457 .
- 4- M. S. El Naschie : “ Coupled oscillations and mode locking of Quantum Gravity fields, Scale Relativity and $\mathcal{E}^{(\infty)}$ space “. Chaos, Solitons and Fractals **12** (2001) 179-192.
- 5- C. Castro : “ Hints of a New Relativity Principle from p -Brane Quantum Mechanics “ Chaos, Solitons and Fractals **11** (2000) 1721
- C. Castro : “ Noncommutative Geometry, Negative Probabilities and Cantorian Fractal Spacetime “ Chaos, Solitons and Fractals **12** (2001) 101-104
- 6- C. Castro, A. Granik : “ Scale Relativity in $\mathcal{E}^{(\infty)}$ space and the Average Dimension of the World ’ To appear in Chaos, Solitons and Fractals. hep-th/0004152
- 7- L. Nottale : “ Fractal Spacetime and Microphysics : Towards a theory of Scale Relativity. World Scientific , 1993.
- L. Nottale : “ La Relativite dans tous ses Etats “ Hachette Literature, Paris 1998.
- 8- J. Argyris, C. Ciubotariu H. Matuttis : “ Fractal space, cosmic strings and spontaneous symmetry breaking “ Chaos, Solitons and Fractals **12** (2001) 1-48